

Congratulations!




You've taken the first step toward self-injection of TEZSPIRE. Now here are a few things we think can help you along the way.

In this kit we've provided resources to support you throughout your injection experience and help you get the most from your treatment. Let's do this—together.

Sincerely,
Team TEZSPIRE

Get inspired by other
TEZSPIRE patients.




TEZSPIRE is an add-on maintenance treatment for people aged 12 and older with severe asthma or inadequately controlled chronic rhinosinusitis with nasal polyps (CRSwNP).
TEZSPIRE is not a rescue medicine.

Injection Log



Choose a different site each time that is at least 1 inch away from where you last injected.



Recommended Injection Sites:

- Outer area of upper arm (caregiver only)
- Lower stomach (patient or caregiver: except into the 2-inch area around the belly button)
- Front of thigh (patient or caregiver)

Do not inject:

- Into the 2-inch (5cm) area around your belly button
- Where the skin is tender, bruised, scaly, or hard
- Into scars or damaged skin
- Through clothing

Day / Month

Injection Site

Day / Month	<input type="checkbox"/> Left Upper Arm	<input type="checkbox"/> Right Upper Arm	<input type="checkbox"/> Stomach	<input type="checkbox"/> Left Thigh	<input type="checkbox"/> Right Thigh
/	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
/	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
/	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
/	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
/	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
/	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
/	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
/	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
/	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
/	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
/	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
/	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
/	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
/	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
/	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

IMPORTANT SAFETY INFORMATION

Do not use TEZSPIRE if you are allergic to tezepelumab-ekko or any of its ingredients.

Do not use to treat sudden breathing problems.

TEZSPIRE may cause serious side effects, including:

- **allergic (hypersensitivity) reactions, including anaphylaxis.** Serious allergic reactions can happen after your TEZSPIRE injection. Allergic reactions can sometimes happen hours or days after you get a dose of TEZSPIRE. Call your healthcare provider or get emergency help right away if you get any of the following symptoms of an allergic reaction:
 - rash
 - breathing problems
 - hives
 - red, itchy, swollen, or inflamed eyes
 - swelling of your face, mouth, and tongue
 - fainting, dizziness, feeling lightheaded

Please read full **Prescribing Information**, including **Patient Information** and **Instructions for Use**.

IMPORTANT SAFETY INFORMATION

Do not use TEZSPIRE if you are allergic to tezepelumab-ekko or any of its ingredients.

Do not use to treat sudden breathing problems.

TEZSPIRE may cause serious side effects, including:

- **allergic (hypersensitivity) reactions, including anaphylaxis.** Serious allergic reactions can happen after your TEZSPIRE injection. Allergic reactions can sometimes happen hours or days after you get a dose of TEZSPIRE. Call your healthcare provider or get emergency help right away if you get any of the following symptoms of an allergic reaction:
 - rash
 - breathing problems
 - hives
 - red, itchy, swollen, or inflamed eyes
 - swelling of your face, mouth, and tongue
 - fainting, dizziness, feeling lightheaded

Before using TEZSPIRE, tell your healthcare provider about all of your medical conditions, including if you:

- have ever had a severe allergic reaction.
- have a parasitic (helminth) infection.
- have recently received or are scheduled to receive any vaccinations. You should not receive a “live vaccine” if you are treated with TEZSPIRE.
- are pregnant or plan to become pregnant.
- are breastfeeding or plan to breastfeed. It is not known if TEZSPIRE passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby if you use TEZSPIRE.
- are taking prescription and over-the-counter medicines, vitamins, or herbal supplements.

Do not change or stop taking your other asthma medicines unless instructed to do so by your healthcare provider.

The most common side effects of TEZSPIRE include: Sore throat, joint and back pain, common cold symptoms, flu, upper

respiratory tract infections, injection site reactions, and nose bleeds. These are not all the possible side effects.

What is TEZSPIRE?

TEZSPIRE is a prescription medicine used:

- with other asthma medicines for the maintenance treatment of severe asthma in people 12 years of age and older whose asthma is not controlled with their current asthma medicine.
- with other medicines for the maintenance treatment of chronic rhinosinusitis with nasal polyps (CRSwNP) in people 12 years of age and older whose CRSwNP is not controlled with their current CRSwNP medicine.

TEZSPIRE helps:

- prevent severe asthma attacks (exacerbations) and can improve your breathing.
- reduce the size of your polyps, improve symptoms including i.e. nasal congestion and loss of smell, and reduce the need for oral corticosteroids and surgery for your nasal polyps in CRSwNP.

TEZSPIRE is not used to treat sudden breathing problems. Tell your healthcare provider if your asthma does not get better or if it gets worse after you start treatment with TEZSPIRE.

You may report side effects related to AstraZeneca products.

Please read full Prescribing Information, including Patient Information and Instructions for Use.

 **TEZSPIRE**[®]
(tezepelumab-ekko) Subcutaneous
Injection 210 mg

AMGEN[®] AstraZeneca 

TEZSPIRE is an add-on maintenance treatment for people aged 12 and older with severe asthma or inadequately controlled chronic rhinosinusitis with nasal polyps (CRSwNP). TEZSPIRE is not a rescue medicine.

Injection Guide



This resource is intended to provide you with an overview and basic information about injecting your medication and does not replace the Instructions for Use. Prior to administering your medication, thoroughly review the complete Instructions for Use and the full Prescribing Information provided with your medication with your doctor. If you have any questions, please contact your doctor.

Before injecting:




Remove pen from refrigerator and allow it to come to room temperature (at least 60 min). Use within 30 days of removal.

Make sure the liquid is colorless to light yellow.




Wash your hands and clean the injection site with an alcohol wipe.

Key injection steps:




POSITION the pen with the needle guard straight toward the skin (90-degree angle).

- ▶ Make sure you can see the viewing window
- ▶ You can either gently pinch the skin at the injection site or give the injection without pinching the skin



PRESS & HOLD the pen down firmly against your skin for about 15 seconds.


- ▶ Make sure you cannot see the orange needle guard anymore
- ▶ You will hear the first “click,” which means the injection has started
- ▶ The orange plunger will move down in the viewing window



CONFIRM the viewing window turns completely orange.

- ▶ You will hear a second “click,” which means the injection has finished
- ▶ Lift the pen straight up after you have completed your injection. The orange needle guard will slide down and lock into place

For supplemental injection information, please watch the demonstration video.



IMPORTANT SAFETY INFORMATION

Do not use TEZSPIRE if you are allergic to tezepelumab-ekko or any of its ingredients.

Do not use to treat sudden breathing problems.

TEZSPIRE may cause serious side effects, including:

- **allergic (hypersensitivity) reactions, including anaphylaxis.** Serious allergic reactions can happen after your TEZSPIRE injection. Allergic reactions can sometimes happen hours or days after you get a dose of TEZSPIRE. Call your healthcare provider or get emergency help right away if you get any of the following symptoms of an allergic reaction:
 - rash
 - breathing problems
 - hives
 - red, itchy, swollen, or inflamed eyes
 - swelling of your face, mouth, and tongue
 - fainting, dizziness, feeling lightheaded

Please read full **Prescribing Information**, including **Patient Information** and **Instructions for Use**.

IMPORTANT SAFETY INFORMATION

Do not use TEZSPIRE if you are allergic to tezepelumab-ekko or any of its ingredients.

Do not use to treat sudden breathing problems.

TEZSPIRE may cause serious side effects, including:

- **allergic (hypersensitivity) reactions, including anaphylaxis.** Serious allergic reactions can happen after your TEZSPIRE injection. Allergic reactions can sometimes happen hours or days after you get a dose of TEZSPIRE. Call your healthcare provider or get emergency help right away if you get any of the following symptoms of an allergic reaction:
 - rash
 - breathing problems
 - hives
 - red, itchy, swollen, or inflamed eyes
 - swelling of your face, mouth, and tongue
 - fainting, dizziness, feeling lightheaded

Before using TEZSPIRE, tell your healthcare provider about all of your medical conditions, including if you:

- have ever had a severe allergic reaction.
- have a parasitic (helminth) infection.
- have recently received or are scheduled to receive any vaccinations. You should not receive a “live vaccine” if you are treated with TEZSPIRE.
- are pregnant or plan to become pregnant.
- are breastfeeding or plan to breastfeed. It is not known if TEZSPIRE passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby if you use TEZSPIRE.
- are taking prescription and over-the-counter medicines, vitamins, or herbal supplements.

Do not change or stop taking your other asthma medicines unless instructed to do so by your healthcare provider.

The most common side effects of TEZSPIRE include: Sore throat, joint and back pain, common cold symptoms, flu, upper

respiratory tract infections, injection site reactions, and nose bleeds. These are not all the possible side effects.

What is TEZSPIRE?

TEZSPIRE is a prescription medicine used:

- with other asthma medicines for the maintenance treatment of severe asthma in people 12 years of age and older whose asthma is not controlled with their current asthma medicine.
- with other medicines for the maintenance treatment of chronic rhinosinusitis with nasal polyps (CRSwNP) in people 12 years of age and older whose CRSwNP is not controlled with their current CRSwNP medicine.

TEZSPIRE helps:

- prevent severe asthma attacks (exacerbations) and can improve your breathing.
- reduce the size of your polyps, improve symptoms including i.e. nasal congestion and loss of smell, and reduce the need for oral corticosteroids and surgery for your nasal polyps in CRSwNP.

TEZSPIRE is not used to treat sudden breathing problems. Tell your healthcare provider if your asthma does not get better or if it gets worse after you start treatment with TEZSPIRE.

You may report side effects related to AstraZeneca products [↗](#).

Please read full Prescribing Information, including Patient Information and Instructions for Use.

 **TEZSPIRE**[®]
(tezepelumab-ekko) Subcutaneous
Injection 210 mg

AMGEN[®] AstraZeneca 

TEZSPIRE IS AN ADD-ON MAINTENANCE TREATMENT FOR PEOPLE AGED 12 AND OLDER WITH SEVERE ASTHMA OR INADEQUATELY CONTROLLED CHRONIC RHINOSINUSITIS WITH NASAL POLYPS (CRSWNP).
TEZSPIRE IS NOT A RESCUE MEDICINE.

Tools to Help You

You can use these resources to help along the way.

 **TEZSPIRE**[®]
(tezepelumab-ekko) Subcutaneous Injection 210 mg




When you get your pen, leave it in its original carton and place it directly into the refrigerator.*

Remove the pen from the refrigerator about 60 minutes before use to allow it to come to room temperature.†

*Store TEZSPIRE in a refrigerator between 36°F to 46°F (2°C to 8°C) in its original carton until you're ready to use it. Store TEZSPIRE in the original carton to protect it from light.

†Do not put TEZSPIRE back in the refrigerator after it has reached room temperature. Throw away (dispose of) TEZSPIRE that has been stored at room temperature for more than 30 days.

My TEZSPIRE Day



Write the day of the month here or add it to your calendar.



Want a little extra help?
Call a nurse educator[‡]
1-888-TZSPIRE

[‡]Nurse educators are nurses by training, but they are not part of your treatment team or an extension of your doctor's office. Nurse educators do not inject patients with TEZSPIRE. You will be referred to your doctor's office for clinical advice.

Eligible commercially insured patients may pay as little as

\$0[§]

Get doses for as little as \$0 and up to \$100 off your in-office administration costs (pre-filled syringe only).

Enroll now



[§]For eligible commercially insured patients only. Eligibility criteria and program maximums apply. Patients who are residents of Massachusetts or Rhode Island are not eligible for injection administration support. Visit TEZSPIRE.COM for full Terms and Conditions.

Important!

- Your specialty pharmacy (SP) will contact you to coordinate shipment
- Add their phone number to your contacts
- Request a free sharps disposal from your SP



IMPORTANT SAFETY INFORMATION

Do not use TEZSPIRE if you are allergic to tezepelumab-ekko or any of its ingredients.

Do not use to treat sudden breathing problems.

TEZSPIRE may cause serious side effects, including:

- **allergic (hypersensitivity) reactions, including anaphylaxis.** Serious allergic reactions can happen after your TEZSPIRE injection. Allergic reactions can sometimes happen hours or days after you get a dose of TEZSPIRE. Call your healthcare provider or get emergency help right away if you get any of the following symptoms of an allergic reaction:
 - rash
 - breathing problems
 - hives
 - red, itchy, swollen, or inflamed eyes
 - swelling of your face, mouth, and tongue
 - fainting, dizziness, feeling lightheaded

Please read full **Prescribing Information**, including **Patient Information** and **Instructions for Use**.

TEZSPIRE for Severe Asthma

71%
FEWER
ASTHMA
ATTACKS
in one trial[¶]

**PREVENT
ASTHMA
ATTACKS**

Another trial saw 56% reduction of asthma attacks compared to patients not taking TEZSPIRE.^{¶¶}

[¶]In one study, TEZSPIRE patients had 0.20 asthma attacks a year vs 0.72 in patients taking placebo.

^{¶¶}In another study, TEZSPIRE patients had 0.93 asthma attacks a year vs 2.10 in patients taking placebo.

**TEZSPIRE CAN HELP
YOU BREATHE
BETTER AND
RELIEVE YOUR
ASTHMA SYMPTOMS**

TEZSPIRE for CRSwNP (Chronic Rhinosinusitis with Nasal Polyps)



TEZSPIRE was proven to help shrink nasal polyps and reduce nasal congestion.



Patients experienced a 92% reduction in the need for surgery and/or steroids after starting TEZSPIRE.

In a clinical study, the proportion of patients needing nasal polyp surgery and/or systemic corticosteroids was 5.7% in the group taking TEZSPIRE, and 30.6% in the group who didn't take TEZSPIRE.

Results may vary.

IMPORTANT SAFETY INFORMATION (continued)

Before using TEZSPIRE, tell your healthcare provider about all of your medical conditions, including if you:

- have ever had a severe allergic reaction.
- have a parasitic (helminth) infection.
- have recently received or are scheduled to receive any vaccinations. You should not receive a "live vaccine" if you are treated with TEZSPIRE.
- are pregnant or plan to become pregnant.

Please read full [Prescribing Information](#), including [Patient Information](#) and [Instructions for Use](#).

IMPORTANT SAFETY INFORMATION

Do not use TEZSPIRE if you are allergic to tezepelumab-ekko or any of its ingredients.

Do not use to treat sudden breathing problems.

TEZSPIRE may cause serious side effects, including:

- **allergic (hypersensitivity) reactions, including anaphylaxis.** Serious allergic reactions can happen after your TEZSPIRE injection. Allergic reactions can sometimes happen hours or days after you get a dose of TEZSPIRE. Call your healthcare provider or get emergency help right away if you get any of the following symptoms of an allergic reaction:
 - rash
 - breathing problems
 - hives
 - red, itchy, swollen, or inflamed eyes
 - swelling of your face, mouth, and tongue
 - fainting, dizziness, feeling lightheaded

Before using TEZSPIRE, tell your healthcare provider about all of your medical conditions, including if you:

- have ever had a severe allergic reaction.
- have a parasitic (helminth) infection.
- have recently received or are scheduled to receive any vaccinations. You should not receive a “live vaccine” if you are treated with TEZSPIRE.
- are pregnant or plan to become pregnant.
- are breastfeeding or plan to breastfeed. It is not known if TEZSPIRE passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby if you use TEZSPIRE.
- are taking prescription and over-the-counter medicines, vitamins, or herbal supplements.

Do not change or stop taking your other asthma medicines unless instructed to do so by your healthcare provider.

The most common side effects of TEZSPIRE include: Sore throat, joint and back pain, common cold symptoms, flu, upper

respiratory tract infections, injection site reactions, and nose bleeds. These are not all the possible side effects.

What is TEZSPIRE?

TEZSPIRE is a prescription medicine used:

- with other asthma medicines for the maintenance treatment of severe asthma in people 12 years of age and older whose asthma is not controlled with their current asthma medicine.
- with other medicines for the maintenance treatment of chronic rhinosinusitis with nasal polyps (CRSwNP) in people 12 years of age and older whose CRSwNP is not controlled with their current CRSwNP medicine.

TEZSPIRE helps:

- prevent severe asthma attacks (exacerbations) and can improve your breathing.
- reduce the size of your polyps, improve symptoms including i.e. nasal congestion and loss of smell, and reduce the need for oral corticosteroids and surgery for your nasal polyps in CRSwNP.

TEZSPIRE is not used to treat sudden breathing problems. Tell your healthcare provider if your asthma does not get better or if it gets worse after you start treatment with TEZSPIRE.

You may report side effects related to AstraZeneca products^{1,2}.

Please read full Prescribing Information, including Patient Information and Instructions for Use.



TEZSPIRE is a registered trademark of Amgen Inc. and AstraZeneca. ©2025 AstraZeneca. All rights reserved. US-107067 Last Updated 12/25